arXiv:quant-ph/9706057v2 28 Jun 1997

Exact solution for generalized pairing

Feng Pan[†] **and J. P. Draayer**

Department of Physics & Astronomy, Louisiana State University,

Baton Rouge, LA 70803-4001

Abstract

An infinite dimensional algebra, which is useful for deriving exact solutions of the generalized pairing problem, is introduced. A formalism for diagonalizing the corresponding Hamiltonian is also proposed. The theory is illustrated with some numerical examples.

PACS numbers:21.60.-n, 21.60.Cs, 02.20.Tw, 03.65.Fd

[†]On leave from the Department of Physics, Liaoning Normal University, Dalian 116029, P. R. China

Pairing has long been considered an important interaction in physics. The concept can be traced back to the seniority scheme introduced by Racah in atomic physics.[1] Its physical significance was first realized in the study of superconductivity.[2] Following the suggestions of Bohr, Mottelson, and Pines,[3] the first detailed application of pairing in nuclei was made by Belyaev.[4] The concept has since been applied to other phenomena: high T_c superconductivity,[5,6] applications using the Hubbard model,[7] pairing phenomena in liquids,[8] and metal clusters.[9]

BCS methods have yielded major successes in studies of superconductivity. When applied to nuclei, however, some negatives come with the positives. First of all, not only is the number of nucleons in a nucleus typically small, the number of valence particles ($n \sim 10$) which dominates the behaviour of low-lying states is too few to support underlying assumptions of the BCS approximation, specifically, $\delta n/n$ is not negligible. As a consequence, particle-number-nonconservation effects enter and can lead to serious difficulties, such as spurious states, nonorthogonal solutions, etc. Secondly, an essential feature of pairing correlations are even-odd differences, which are driven mainly by Pauli blocking. It is difficult to treat these differences in the BCS formalism because different quasi-particle bases must be introduced for different blocked levels. For these reasons, a particle-number-conserving method for treating the pairing problem has been suggested for well-deformed nuclei.[10] The method uses a configuration energy truncation scheme, and takes the strength of the pairing interaction to be the same for all orbitals. In this limit the pairing Hamiltonian can be diagonalized in truncated configuration spaces. Because the theory applies to well-deformed nuclei, each orbital can only be occupied by a single pair of particles.

The generalized pairing Hamiltonian for spherical nuclei can be written as

$$\hat{H} = \sum_{jm} \epsilon_j a^\dagger_{jm} a_{jm} - |G| \sum_{jj'} c_{jj'} S^+(j) S^-(j), \tag{1}$$

where the ϵ_j are single-particle energies and $S^\pm(j)$ and $S^0(j)$ are the pairing operators for a single-j shell defined by

$$\begin{aligned} S^+(j) &= \sum_{m>0} (-)^{j-m} a^\dagger_{jm} a^\dagger_{j-m}, \\ S^-(j) &= \sum_{m>0} (-)^{j-m} a_{j-m} a_{jm}, \\ S^0(j) = \frac{1}{2} \sum_{m>0} (a^\dagger_{jm} a_{jm} + a^\dagger_{j-m} a_{j-m} - 1) &= \frac{1}{2}(\hat{N}_j - \Omega_j), \end{aligned} \tag{2}$$

where $\Omega_j \equiv j + 1/2$ is the maximum number of pairs in the j-th shell, $\hat{N}_j$ is the particle number operator for the j-th shell, and the $c_{jj'}$ measure the pairing strength between

different j-shells. In general, for N pairs, Hamiltonian (1) can be diagonalized in bases states that are products of the single-j shell pairing wave functions:

$$|N>=\sum_{k_i} B_{k_1 k_2 \cdots k_p} S_{j_1}^{+\ k_1} S_{j_2}^{+\ k_2} S_{j_3}^{+\ k_3} \cdots S_{j_p}^{+\ k_p} |0>, \tag{3}$$

where the summation is restricted by

$$\sum_{i=1}^{p} k_i = N, \tag{4}$$

the $B_{k_1 k_2 \cdots k_p}$ are expansion coefficient that need to be determined, and $|0>$ is the pairing vacuum state which satisfies the condition

$$S_j^- |0> \ = \ 0 \quad \text{for all} \ \ j. \tag{5}$$

The dimensionality of the Hamiltonian matrix in this basis increases very rapidly with increasing N and the number of shells p. It is less than or equal to the dimension of the irreducible representation (irrep) $[N\dot{0}]$ of the unitary group $U(p)$ due to the Pauli principle,

$$\dim \ \leq \ \frac{(p+N-1)!}{N!(p-1)!}. \tag{6}$$

The equal sign holds in (6) when all the single-j shell pairing wave functions in the summation of (3) are Pauli allowed. Indeed, the problem quickly becomes intractable because there are no analytical expressions or recursion relations for determining the $B_{k_1 k_2 \cdots k_p}$ coefficients.

Following the quasi-spin approximation,[11] consider a simpler Hamiltonian

$$\hat{H} = -|G| S_0^+ S_0^-, \tag{7}$$

where

$$S_0^+ = \sum_j c_j^* S^+(j), \ S_0^- = \sum_j c_j S^-(j) \tag{8}$$

with the coefficient $c_i, \ c_j^*$ satisfying the condition

$$\sum_i |c_i|^2 = 1. \tag{9}$$

This defines generalized pairing as proposed by Talmi.[12] Clearly, in the notation of (1), $c_{jj'} = c_j^* c_{j'}$ with $|c_j|^2$ giving the percentage of single-j shell pairing in the Hamiltonian. In what follows, the c_j are taken to be real.

To diagonalize Hamiltonian (7), consider an algebra generated by

$$S_m^0 = \sum_j c_j^{2m} S^0(j),$$

$$S_m^{\pm} = \sum_j c_j^{2m+1} S^{\pm}(j). \tag{10}$$

It is easy to show that these generators satisfy the following commutation relations:

$$[S_m^+,\ S_n^-] \ = \ 2S_{m+n+1}^0,$$

$$[S_m^0,\ S_n^{\pm}] \ = \ \pm\, S_{m+n}^{\pm}. \tag{11}$$

Therefore, the $\{S_m^\mu,\ \mu = 0, +, -;\ m = 0, \pm 1, \pm 2, \cdots\}$ form an infinite-dimensional algebra, one that differs only slightly from a general Lie algebra of the affine type.

The unique lowest-weight state of this algebra is simply the product of the single-j shell pairing vacua with arbitrary seniority quantum numbers. Therefore, it suffices to consider the total seniority zero case. The lowest-weight state satisfies

$$S_m^-|0>= 0;\ m = 0,\ \pm 1,\ \pm 2,\ \cdots, \tag{12}$$

and

$$S_m^0|0>= -\frac{1}{2}\sum_j |c_j|^{2m}\Omega_j|0>= \Lambda_m|0> . \tag{13}$$

Furthermore, it can be proven that the eigenvectors of $\hat{H}$ for any N and non-zero energy eigenvalue can be expanded as

$$|N>= \sum_{n_i} x_1^{n_1} x_2^{n_2} \cdots x_{N-1}^{n_{N-1}} S_0^+ S_{n_1}^+ S_{n_2}^+ \cdots S_{n_{N-1}}^+ |0>, \tag{14}$$

where

$$n_i = \begin{cases} -1, \ -2, \ \cdots \ \text{if} \ \ c_j^2 x_i \ > 1, \\ \\ 0, \ 1, \ 2, \ \cdots \ \text{if} \ \ c_j^2 x_i \ < 1. \end{cases} \tag{15}$$

In any case, up to a normalization constant (14) can always be written as

$$|N >= S_0^+ S_{x_1}^+ S_{x_2}^+ \cdots S_{x_{N-1}}^+ |0 >, \tag{16}$$

where

$$S_{x_i}^+ = \sum_j \frac{c_j}{1 - c_j^2 x_i} S_j^+. \tag{17}$$

While $x_1, \ x_2, \ \cdots, x_{N-1}$ are real or complex numbers satisfying the relation

$$-\frac{1}{2} \sum_{j=1}^{p} \Omega_j c_j^2 \alpha \frac{1}{1 - \alpha y_i c_j^2} = \frac{1}{y_i} + \sum_{k \neq i} \frac{1}{y_i - y_k}, \quad i = 1, \ 2, \cdots, \ N-1, \tag{18}$$

with

$$\sum_{i=1}^{N-1} \frac{1}{y_i} = 1, \tag{19}$$

where

$$y_i = x_i / \alpha, \quad \alpha = -\frac{2}{h + 2\Lambda_1}, \quad h \equiv E/(-|G|). \tag{20}$$

Therefore, the coefficients x_i (i=1, 2,$\cdots$, N-1) and eigenvalues of the pairing energy $E \neq 0$ are simultaneously determined by the system of equations (18) and (19).

While eigenvectors for $E = 0$ can be expanded as follows

$$|N >= \sum_{n_i} x_1^{n_1} x_2^{n_2} \cdots x_N^{n_N} S_{n_1}^+ S_{n_2}^+ \cdots S_{n_N}^+ |0 >, \tag{21}$$

where the restrictions on the integers n_i are the same as those given by (15), and the expansion coefficients x_i are determined by the following set of equations

$$\sum_j \Omega_j \frac{c_j^2}{1 - x_1 c_j^2} = 0 \quad \text{for} \ \ N = 1, \tag{22}$$

$$\sum_j \Omega_j \frac{c_j^2}{1 - x_i c_j^2} = \sum_{k \neq i} \frac{1}{x_i - x_k}, \quad i = 1,\ 2,\ \cdots,\ N, \quad \text{for} \quad N \geq 2. \tag{23}$$

It is instructive to write down the first few energy eigenvalues and eigenstates.

$$\hat{h}|0>= 0; \tag{24}$$

$$\hat{h}|1>= \sum_j |c_j|^2 \Omega_j |1>, \quad |1>= S_0^+|0>,$$

$$\hat{h}|N=1,\ \rho> \ = \ 0, \qquad |N=1,\ \rho>= \sum_j \frac{c_j}{1 - x^\rho c_j^2} S_j^+|0>, \quad \rho \ = \ 1,\ 2,\ \cdots, \tag{25}$$

where x_ρ is determined by (22);

$$\hat{h}|2>= \ h|2>, \quad |2>= S_0^+ S_x^+|0>, \tag{26}$$

where h is one of the solutions of the equation

$$\frac{1}{2} \sum_{j=1}^{p} \Omega_j c_j^2 x \frac{1}{x c_j^2 - 1} \ = \ 1 \tag{27}$$

where

$$x = -\frac{2}{h + 2\Lambda_1}, \tag{28}$$

$$\hat{h}|N=2,\ \rho \ > \ = 0, \quad \rho \ = \ 1,\ 2,\ \cdots \tag{29}$$

where

$$|N=2,\ \rho>= \sum_{j\ j'} \frac{c_j c_{j'}}{(1 - x_1^\rho c_j^2)(1 - x_2^\rho c_{j'}^2)} S_j^+ S_{j'}^+|0>, \tag{30}$$

and x_i^ρ are determined by (23).

Energy levels of the generalized pairing interaction for the $j \ = \ 1/2,\ 3/2$, and $5/2$ case with $c_{1/2} = \sqrt{0.1}$, $c_{3/2} = \sqrt{0.3}$, and $c_{5/2} \ = \ \sqrt{0.6}$ are shown in Fig. 1. It can be seen from Table 1 that the energy level with the largest c_j value for the highest j orbit is the lowest one for any fixed N. Finally, an example of the lowest levels for given N in the 5-th and 6-th shell, respectively, are shown in Table 2.

Acknowledgment

Supported by the National Science Foundation through Cooperative Agreement No. EPS-9550481 and Grant No. 96030006.

---

[1] G. Racah, Phys. Rev. **62**, 438 (1942)
[2] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **108**, 1175 (1957)
[3] A. Bohr, B. R. Mottelson, and D. Pines, Phys. Rev. **110**, 936 (1958)
[4] S. T. Belyaev, Mat. Fys. Medd. **31**, 11 (1959)
[5] M. Randeria, J. M. Duan, and L. Y. Shieh, Phys. Rev. Lett. **62**, 981 (1989)
[6] S. Schmitt-Rink, C. M. Verma, and A. E. Ruckenstein, Phys. Rev. Lett. **63**, 445 (1989)
[7] C. N. Yang, Phys. Rev. Lett. **63**, 2144 (1989)
[8] D. W. Cooper, J. S. Batchelder, and M. A. Taubenblatt, J. Coll. Int. Sci. **144**, 201 (1991)
[9] M. Barranco, S. Hernandez, and R. J. Lombard, Z. Phys. **D22**, 659(1992)
[10] J. Y. Zeng and T. S. Cheng, Nucl. Phys. **A405**, 1 (1983); **A411**, 49 (1984); **A414**, 253 (1984)
[11] A. K. Kerman, Ann. Phys. (N. Y.) **12**, 300 (1961)
[12] I. Talmi, Nucl. Phys. **A172**, 1 (1971)

Table 1. The lowest energy levels for $j = 1/2,\ 3/2,\ 5/2$ with:

Case a: $c_{1/2} = \sqrt{0.1},\ c_{3/2} = \sqrt{0.2},\ c_{5/2} = \sqrt{0.7}$;
Case b: $c_{1/2} = \sqrt{0.2},\ c_{3/2} = \sqrt{0.1},\ c_{5/2} = \sqrt{0.7}$;
Case c: $c_{1/2} = \sqrt{0.2},\ c_{3/2} = \sqrt{0.7},\ c_{5/2} = \sqrt{0.1}$;
Case d: $c_{1/2} = \sqrt{0.7},\ c_{3/2} = \sqrt{0.2},\ c_{5/2} = \sqrt{0.1}$.

	$h = E/(-\|G\|)$			
N	a	b	c	d
0	0	0	0	0
1	2.6	2.5	1.9	1.4
2	4.083	3.866	2.858	2.136
3	4.735	4.441	3.278	2.481
4	4.674	4.377	3.100	2.320
5	4.084	3.866	2.858	2.136
6	2.600	2.500	1.900	1.400

Table 2. Lowest energy levels for 5th and 6th shells. The parameters are chosen as follows: $c_{1/2} = \sqrt{0.2},\ c_{3/2} = \sqrt{0.3},\ c_{5/2} = \sqrt{0.22},\ c_{9/2} = \sqrt{0.28}$ for 5th shell, whereas $c_{1/2} = \sqrt{0.2}$, $c_{3/2} = \sqrt{0.14},\ c_{5/2} = \sqrt{0.22},\ c_{7/2} = \sqrt{0.28},\ c_{11/2} = \sqrt{0.16}$ for 6th shell, respectively.

	$(N,\ h = E/(-\|G\|)\ )$					
5th shell	(0, 0)	(1, 2.860)	(2, 5.192)	(3, 6.999)	(4, 8.283)	(5, 9.046)
	(6, 9.292)	(7, 9.024)	(8, 8.243)	(9, 6.954)	(10, 5.515)	(11, 2.86)
6th shell	(0,0)	(1, 3.220)	(2, 6.014)	(3, 8.387)	(4, 10.345)	(5, 11.893)
	(6, 13.035)	(7, 13.778)	(8, 14.125)	(9, 14.082)	(10, 13.653)	(11, 12.843)
	(12, 11.655)	(13, 10.096)	(14, 8.167)	(15, 5.874)	(16, 3.220)	



Fig. 1. An example of excited energy levels for the generalized pairing interaction with $N \leq 3$, where the number on the left indicates the degeneracy when it is greater than one.